

Table 2. Date of birth and BMD values (gr/cm²) in the area of the femoral neck during the two years of study, for the experimental group.

Date of birth	BMD (gr/cm ²)		
	0	12 months	24 months
13/7/1989	1,177	1,188	1,204
13/5/1987	1,135	1,185	1,273
12/6/1987	0,940	0,949	0,973
12/12/1987	0,943	0,954	0,976
15/12/1986	0,948	0,958	1,237
12/3/1988	1,203	1,215	1,235
2/1/1987	0,805	0,805	0,836
26/7/1986	0,833	0,840	0,858
30/6/1987	0,945	1,017	1,039
31/3/1988	1,009	0,955	0,972
5/8/1987	1,029	1,101	1,685
31/8/1987	0,990	1,000	1,019
21/8/1987	0,982	0,99	1,020
4/9/1987	0,977	0,99	1,022
8/9/1988	0,993	1,001	1,025
2/10/1989	0,952	1,003	1,024
8/8/1989	1,002	1,009	1,021
23/12/1986	0,971	0,983	1,001
6/9/1988	0,969	0,982	1,002
31/7/1988	0,980	0,980	1,005
5/5/1987	0,978	0,981	0,998
23/8/1989	0,992	0,986	1,019
12/1/1988	0,983	1,002	1,022
22/7/1987	0,984	0,99	1,009
23/8/1989	0,971	1,002	1,008
20/1/1988	1,001	1,009	1,031
5/4/1988	0,988	0,985	1,005
12/3/1988	1,203	1,100	1,237
13/5/1987	1,193	1,185	1,229
5/6/1988	0,987	1,005	1,025