

Table 4. Date of birth and BMD values (gr/cm²) in the area of the femoral neck during the two years of study for the control group.

Date of birth	BMD (gr/cm ²)		
	0	12 months	24 months
12/3/1988	1,203	1,211	1,237
13/5/1989	1,085	1,190	1,204
12/5/1986	1,183	1,192	1,090
12/5/1986	1,183	1,192	1,218
13/4/1988	1,102	1,173	1,229
2/9/1987	0,838	0,868	0,893
2/2/1989	0,920	0,944	0,988
22/11/1988	0,970	0,987	0,999
13/5/1987	0,990	1,00	1,011
14/6/1986	1,09	1,104	1,123
15/8/1987	0,99	0,981	1,017
14/2/1986	0,960	0,968	0,987
9/9/1990	1,001	1,010	1,031
4/5/1988	0,966	0,977	0,992
7/11/1986	0,961	0,975	0,991
18/11/1988	0,950	0,943	0,978
23/1/1987	0,960	0,977	0,992
1/8/1985	0,97	0,980	0,999
22/10/1989	0,98	0,988	0,989
11/11/1988	0,988	0,999	1,099
3/2/1988	0,944	0,955	0,971
9/8/1988	0,90	0,976	0,991
17/6/1986	1,003	1,015	1,048
1/8/1986	1,101	1,170	1,222
25/4/1989	0,949	0,944	0,976
29/3/1988	0,989	1,102	0,977
31/3/1989	0,949	0,941	0,977
14/12/1988	1,098	1,100	1,137
19/4/1988	0,947	0,942	0,976
22/4/1988	0,968	1,099	1,016
8/5/1988	1,000	1,009	1,029
18/4/1990	1,002	1,009	1,035
19/12/1986	1,083	1,189	1,201
25/7/1988	0,977	0,998	1,005
29/3/1988	0,989	1,102	0,977
8/9/1988	0,970	0,980	1,002
16/7/1989	0,993	1,022	1,032