**Supplementary** Table 1. Maternal and offspring characteristics in study population. MATERNAL CHARACTERISTICS AND OFFSPRING CHARACTERISTICS MEAN (SD) / N (%) N (%) MISSING Age in years, mean (sd) 31.53 (4.43) 0(0)Height in cm, mean (sd) 165.20 (5.35) 0(0)Weight before pregnancy in kg, mean (sd) 62.35 (11.38) 0(0)BMI before pregnancy, mean (sd) 22.81 (3.75) 0(0)Weight at term in kg, mean (sd) 75.81 (12.63) 0(0)13.46 (6.94) Gestational weight gain in kg, mean (sd) 0(0)Frequency of weight checking in pregnancy, n (%) 1 (0.5) 0(0)Every month 97 (50.3) 71 (36.8) Every week Other 24 (12.4) Number of previous pregnancies, n (%) 0 124 (64.2) 0(0)1 43 (22.3) 2 18 (9.3) 3 6(3.1)4 2 (1.0) 143 (74.1) Number of live children, n (%) 0 0(0)45 (23.3) 1 2 5 (2.6) Number of previous vaginal labors, n (%) 0 155 (80.3) 0(0)1 34 (17.6) 2 4 (2.1) 0 174 (90.2) 0(0)Number of previous caesarian sections, n (%) 1 19 (9.8) Use of IVF, n (%) No 184 (95.3) 0(0)Yes 9 (4.7) 187 (96.9) Multiple Pregnancies, n (%) No 0(0)Yes 6(3.1)Smoking before pregnancy, n (%) No 108 (56.0) 0(0)Yes 85 (44.0) Duration of smoking before pregnancy in years1, mean (sd) 4.70 (6.17) 0(0)18 (21.4) Amount of smoking per day before pregnancy2, n (%) 5 cigarettes 1(1.2)10 cigarettes 34 (40.5) 20 cigarettes 26 (31.0) 30 or more cigarettes 6(7.1)88 (45.6) 0(0)Self-reported physical activity before pregnancy, n (%) Low Moderate 76 (39.4)

> Vigorous Professional

No

Yes

No

Yes

No Yes

No

Yes

No

Yes

No

Yes

No

Yes

None

Few days
One month

Negative Probably negative

Neutral

Positive

More than one month

Probably positive

Cervical mucus

Uterine contractions

Caesarian section

Rupture of

membranes
Labor induction

Vaginal birth

Female

Female

Male

No

Yes

Nο

Yes

No

Yes

Male

None

5 cigarettes

10 cigarettes

20 cigarettes

30 or more cigarettes

Smoking during pregnancy, n (%)

Amount of smoking per day during pregnancy, n (%)

Vomiting episodes during pregnancy, n (%)

Bed rest due to vaginal bleeding, n (%)

Bed rest due to placenta abruption, n (%)

Other reasons for bed rest, n (%)

Duration of bedrest, n (%)

Onset of labor, n (%)

Mode of birth, n (%)

Sex, n (%)

Birth weight in g, mean (sd)

Sex of second twin3, n (%)

Total sample size (n)

Birth weight of second twin3, mean (sd)

Willingness to exercise after birth, n (%)

Willingness to exclusively breastfeed, n (%)

Willingness to quit smoking after birth3, n (%)

<sup>1</sup>Including non-smokers, <sup>2</sup>only among smokers, <sup>3</sup>among multiple pregnancies

Abbreviations: sd standard deviation, n number of women, cm centimeters, kg kilograms, g grams

n (%)

Medically indicated bed rest during pregnancy, n (%)

Gynecologist opinion for physical activity during pregnancy,

Gestational age at birth in completed weeks, mean (sd)

Exposure to passive smoking at home during pregnancy, n (%)

26 (13.5)

3(1.6)

170 (88.1)

23 (11.9) 170 (88.1)

17 (8.8)

4 (2.1)

1(0.5)

1(0.5)

105 (54.4)

88 (45.6)

97 (50.3)

96 (49.7)

158 (81.9)

35 (18.1)

175 (90.7)

18 (9.3)

182 (94.3)

11 (5.7)

181 (93.8)

12 (6.2) 156 (80.8)

15 (7.8)

10 (5.2)

12 (6.2) 13 (6.7)

12 (6.2)

106 (54.9)

26 (13.5)

36 (18.7) 38.53 (1.38)

6 (3.2)

29 (15.3)

117 (61.9)

37 (19.6)

92 (47.9)

100 (52.1)

3075.12 (499.66)

88 (52.4)

80 (47.6)

2192.00 (258.79)

5 (100.0)

0(0.0)

104 (53.9)

89 (46.1)

22 (11.4)

171 (88.6)

8 (10.7)

67 (89.3)

193

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

0(0)

4 (2.1)

1 (0.5)

26 (13)

25 (13)

1 (16.7)

1 (16.7)

0(0)

0(0)

10 (11.8)