

Table 5. Logistic regression of the predictive value of serum cystatin C and free beta hCG for diagnosing preeclampsia as compared to normotensive participants.

VARIABLE	UNIVARIABLE			MULTIVARIABLE		
	ODDS RATIO	95% CONFIDENCE INTERVAL	P-VALUE	ADJUSTED ODDS RATIO	95% CONFIDENCE INTERVAL	P-VALUE
<b><i>HCG</i></b>						
<235iu/l	1.00	Reference	Reference	1.00	Reference	Reference
≥235iu/l	3.20	9.08- 11.26	< 0.001	2.92	4.87 - 6.85	< 0.001
<b><i>Cystatin</i></b>						
<1.8mg/dl	1.00	Reference	Reference	1.00	Reference	Reference
≥1.8mg/dl	2.05	1.04- 4.01	< 0.001	6.84	2.30 20.36	0.001
<b><i>Age (years)</i></b>	1.03	0.98 - 1.07	0.184	0.84	0.68 - 1.04	0.111
<b><i>Body Mass Index(kg/m<sup>2</sup>)</i></b>						
Normal	1.00	Reference	Reference	1.00	Reference	Reference
Overweight	2.9	0.47-17.33	0.254	3.35	0.39-41.26	0.287
Obese	1.4	0.23-9.97	0.786	1.84	0.18-18.56	0.767