

Table 1. Causative factors of OVT

ETIOLOGY	NO* CASES
<i>Pregnancy related**</i>	155
Vaginal delivery	80
Caesarian section	48
Urgent delivery - preterm birth	13
Multiple pregnancy (twins)	17
HELLP	3
<i>Endometrial lesion</i>	
Fibroids	11
IUD	4
Endometriosis	1
Cyst	2
Prolapse	1
<i>Pelvic operations /procedures</i>	
Hysterectomy	11
Salpingectomy	3
Other	4
Hysteroscopy	1
<i>Intraabdominal inflammation</i>	
Hydronephrosis	17
Pyelonephritis	4
Other nephropathy	3
Behcet	3
IBD	2
Gallstones - ERCP	4
<i>Thrombophilia - Hypercoagulable state</i>	
OCP	4
Thrombotic history	4
Venous thromboembolism	1
Stroke	5
Positive Family history	1
APS	2
Lupus anticoagulant	1
APS triple positive	5
Recurrent abortions possible APS	3
Inherited Thrombophilia Factors	2
FV-Leiden	6
Low protein C and/or S	
Hyperhomocysteinemia	
Multiple factors	
<i>COVID-19 infection</i>	7
<i>Idiopathic</i>	19

* Number of reported cases, ** 8 cases antepartum

APS: Antiphospholipid Syndrome, ERCP: Endoscopic Retrograde Cholangiopancreatography, HELLP: Hemolysis, Elevated Liver Enzymes and Low Platelets, IBD: Inflammatory Bowel Disease, IUD: Intrauterine Device, OCP: Oral contraceptive pills,

OVT: Ovarian vein thrombosis