

Table 6. Sensitivity pattern of pregnant participant' urinary isolates.

DRUG	KLEBSIELLA		E COLI		PSEUDOMONAS		ENTEROCOCCUS		CITROBACTER		MORGANELLA		OVERALL % SENSITIVITY
	NO AND		NO AND %		NO AND %		NO AND %		NO AND %		NO AND %		
	%SENSITIVITY		SENSITIVITY		SENSITIVITY		SENSITIVITY		SENSITIVITY		SENSITIVITY		
Amoxicillin	13	7.6	6	0.0	4	0.0	3	0.0	1	0.0	1	0.0	3.6
Co-Amoxicillin	13	46.0	6	33.3	4	0.0	3	33.3	1	0.0	1	100	35.7
Nitrofurantoin	13	61.5	6	100.0	4	0.0	3	33.3	1	0.0	1	100	57.1
Nalidixic	13	61.5	6	83.3	4	0.0	3	33.3	1	0.0	1	100	53.6
Trimethoprim	13	7.6	6	16.7	4	0.0	3	0.0	1	0.0	1	100	10.7
Gentamycin	13	61.5	6	66.7	4	75.0	3	66.7	1	100	1	100	67.9
Ofloxacin	13	92.3	6	100.0	4	75.0	3	66.7	1	100	1	100	89.3
Cefuroxime	13	38.4	6	100.0	4	0.0	3	66.7	1	0.0	1	100	50.0
Cefalexin/Cefalor	13	23.1	6	16.7	4	0.0	3	33.3	1	0.0	1	100	21.4
Ceftazidim	13	92.3	6	100.0	4	100	3	66.7	1	100	1	100	92.9
Cloxacillin	13	7.6	6	0.0	4	0.0	3	0.0	1	0.0	1	0.0	3.6