

Table 1. Basal demographic and clinical characteristics in the two study groups.

| Items | Measure | TAP block (N=50) | Control (N=50) | P-value |
|------------------------------|--------------|---------------------|-------------------|---------|
| Age (years) | Mean±SD | 32.1±3.0 | 31.6±4.0 | ^0.536 |
| | Range | 24.0–39.0 | 21.0–39.0 | |
| BMI (kg/m ²) | Mean±SD | 28.7±2.5 | 29.4±2.5 | ^0.154 |
| | Range | 21.7–36.3 | 23.0–34.2 | |
| Parity,(n, %) | primigravida | 18 (36.0%) | 19 (38.0%) | #0.836 |
| | Multigravida | 32 (64.0%) | 31 (62.0%) | |
| Indications,(n, %) | Repeated CS | 19 (38.0%) | 15 (30.0%) | §0.697 |
| | Postdate | 19 (38.0%) | 20 (40.0%) | |
| | PROM | 7 (14.0%) | 11 (22.0%) | |
| | IUGR | 5 (10.0%) | 4 (8.0%) | |
| Gestational age (Week) | Mean±SD | 39.7±1.4 | 39.9±1.3 | ^0.554 |
| | Range | 37.0–42.0 | 37.0–42.0 | |
| Operation duration (minutes) | Mean±SD | 44.8±4.9 | 46.3±5.8 | ^0.167 |
| | Range | 35.0–54.0 | 36.0–64.0 | |