

Table 2. Patients' pain perception (VAS-10) during rest and during movement among the studied groups.

Time	Measures	TAP block (N=50)	Control (N=50)	^p-value	Relative effect Mean±SE 95% CI
<b>Patients' pain perception (VAS-10) during rest</b>					
Hour-2	Mean±SD	0.1±0.3	0.2±0.4	0.063	-0.1±0.1
	Range	0.0-1.0	0.0-1.0		-0.3-0.0
Hour-4	Mean±SD	0.1±0.3	1.8±1.3	<0.001*	-1.7±0.2
	Range	0.0-1.0	1.0-6.0		-2.0--1.3
Hour-8	Mean±SD	2.0±0.8	4.3±1.0	<0.001*	-2.3±0.2
	Range	1.0-4.0	2.0-6.0		-2.7--2.0
Hour-12	Mean±SD	2.3±0.5	4.0±0.7	<0.001*	-1.7±0.1
	Range	2.0-4.0	3.0-5.0		-1.9--1.5
Hour-24	Mean±SD	1.1±0.5	1.3±0.6	0.070	-0.2±0.1
	Range	1.0-3.0	1.0-3.0		-0.4-0.0
<b>Patients' pain perception (VAS-10) during movement</b>					
Hour-2	Mean±SD	0.3±0.5	0.5±0.5	0.133	-0.2±0.1
	Range	0.0-2.0	0.0-2.0		-0.4-0.0
Hour-4	Mean±SD	0.5±0.7	2.5±0.9	<0.001*	-2.0±0.2
	Range	0.0-3.0	2.0-5.0		-2.3--1.7
Hour-8	Mean±SD	3.1±0.7	5.4±0.9	<0.001*	-2.3±0.2
	Range	2.0-5.0	3.0-7.0		-2.6--2.0
Hour-12	Mean±SD	3.2±0.5	4.5±0.6	<0.001*	-1.4±0.1
	Range	2.0-4.0	3.0-6.0		-1.6--1.1
Hour-24	Mean±SD	1.9±0.4	2.1±0.5	0.089	-0.2±0.1
	Range	1.0-3.0	1.0-4.0		-0.3-0.0