

Table 1. Characteristics of the study groups (n=120)

Indicators	Main group, n=60	Comparison group, n=60	
Age (years)	31.6±1.8	24.4±1.2	p<0.05
Family history of genital prolapse	23 (38.3 %)	9 (15.0 %)	$\chi^2=7.20$, p<0.01
Connective tissue dysplasia	26 (43.3 %)	8 (13.3 %)	$\chi^2=11.86$, p<0.001
Asthenic body type	23 (38.3 %)	11 (18.3 %)	$\chi^2=4.97$, p<0.03
BMI > 25,0 kg/m ²	19 (31.7 %)	7 (11.7 %)	$\chi^2=5.94$, p<0.01
GI diseases	16 (26.7 %)	5 (8.3 %)	$\chi^2=5.77$, p<0.02
Pulmonary diseases	14 (23.3 %)	4 (6.7 %)	$\chi^2=5.29$, p<0.02
Varicose veins	18 (30.0 %)	5 (8.3 %)	$\chi^2=7.75$, p<0.005
Instrumental delivery (vacuum extraction)	11 (18.3 %)	1 (1.7 %)	$\chi^2=7.5$, p<0.006
Fetal weight >3600 g	25 (41.7 %)	9 (15.0 %)	$\chi^2=9.23$, p<0.002
Perineal injury	37 (61.7 %)	11 (18.3 %)	$\chi^2=21.7$, p<0.001
Epidural anaesthesia	28 (46.7 %)	15 (25.0 %)	$\chi^2=5.22$, p<0.02
Prolonged labour	14 (23.3 %)	5 (8.3 %)	$\chi^2=4.0$, p<0.05