

Table 2. Nurses' attitudes toward postpartum pain assessment (no = 20).

		Educational Package				Chi-square	P Value
		Pre		Post			
		No	%	No	%		
During the postpartum period, a mother may shout for many reasons, not from postpartum pain	Disagree	11	55.0%	0	0.0%	15.172	p=0.000*
	Agree	9	45.0%	20	100.0%		
Pain should be assessed before and after pain relief measures	Disagree	8	40.0%	0	0.0%	10.000	p<0.002*
	Agree	12	60.0%	20	100.0%		
Every mother experiences postpartum pain differently	Disagree	10	50.0%	1	5.0%	10.157	p<0.001*
	Agree	10	50.0%	19	95.0%		
The mother should be advised to use pain medication along with pain relief measures	Disagree	13	65.0%	5	25.0%	6.465	p<0.011*
	Agree	7	35.0%	15	75.0%		
Understanding postpartum pain may help manage it better	Disagree	6	30.0%	0	0.0%	7.059	p<0.008*
	Agree	14	70.0%	20	100.0%		
Vital signs are always reliable indicators of the severity of postpartum pain	Disagree	12	60.0%	2	10.0%	10.989	p<0.001*
	Agree	8	40.0%	18	90.0%		

\* The Chi-square statistic is significant at the .05 level.