

Table 4. Attitude of nurses regarding the effect of pain-relief methods during postpartum period (no=20).

		Educational Package				Chi-square	P Value
		Pre		Post			
		No	%	No	%		
Mother should expect pain during postpartum	Disagree	6	30.0%	0	0.0%	7.059	p<0.008*
	Agree	14	70.0%	20	100.0%		
Belief that pain relief in postpartum is necessary	Disagree	7	35.0%	0	0.0%	8.485	p<0.004*
	Agree	13	65.0%	20	100.0%		
Use of methods will influence the progress of postpartum	Disagree	14	70.0%	2	10.0%	15.000	p<0.000*
	Agree	6	30.0%	18	90.0%		
Use of methods for pain relief during postpartum is safer	Disagree	10	50.0%	0	0.0%	13.333	p<0.000*
	Agree	10	50.0%	20	100.0%		
Use of pain-relief methods will increase comfort of mother	Disagree	8	40.0%	0	0.0%	10.000	p<0.002*
	Agree	12	60.0%	20	100.0%		
Opinion regarding the effect of pain relief measures on the ability of mother to cope with pain	Disagree	16	80.0%	1	5.0%	23.018	p<0.000*
	Agree	4	20.0%	19	95.0%		
Do you have any concerns about the use of pain relief measures during the postpartum period?	Disagree	14	70.0%	0	0.0%	21.538	p<0.000*
	Agree	6	30.0%	20	100.0%		