

Table 4. Obstetric complications stratified by BMI category with adjusted odds ratios.

<b>Complication</b>	<b>Normal BMI (n=261)</b>	<b>Overweight (n=100)</b>	<b>Obesity (n=160)</b>	<b>Overweight vs Normal BMI, OR (95% CI)</b>	<b>Obesity vs Normal BMI, OR (95% CI)</b>
<b>Gestational hypertension</b>	6 (2.3%)	10 (10.0%)	33 (20.6%)	4.8 (1.7-13.3)*	11.2 (4.6-27.2)*
<b>GDM</b>	14 (5.3%)	6 (6.0%)	13 (8.1%)	1.1 (0.4-3.1)	1.6 (0.7-3.5)
<b>SGA</b>	20 (7.6%)	6 (6.0%)	10 (6.3%)	0.8 (0.3-2.0)	0.8 (0.4-1.7)
<b>Macrosomia</b>	7 (2.6%)	3 (3.0%)	8 (5.0%)	1.1 (0.3-4.3)	2.0 (0.7-5.5)
<b>Preeclampsia with severe features</b>	2 (0.8%)	2 (2.0%)	8 (5.0%)	2.6 (0.4-18.4)	6.7 (1.4-31.6)*
<b>Preeclampsia</b>	3 (1.1%)	1 (1.0%)	3 (1.9%)	0.9 (0.1-8.6)	1.6 (0.3-8.4)
<b>Cesarean section</b>	59 (22.6%)	28 (28.0%)	59 (36.9%)	1.3 (0.8-2.3)	2.0 (1.3-3.1)*
<b>Any complication</b>	90 (34.5%)	42 (42.0%)	88 (55.0%)	1.4 (0.9-2.2)	2.3 (1.6-3.4)*

\*Statistically significant (p&lt;0.05)